

Some remarks on n-fibrations

Lab Lunch, Birmingham

Sina Hazratpour

`sinhp@github.io`

January 2018

Outline of the talk

- In the first section I will review essentials of theory of Grothendieck and Street fibrations of categories. They will be the base case of inductive definition of n -fibration in the last section.

Outline of the talk

- In the first section I will review essentials of theory of Grothendieck and Street fibrations of categories. They will be the base case of inductive definition of n -fibration in the last section.
- In the second section I will mention how to generalize from Grothendieck fibration to 2-fibration of 2-categories and bicategories.

Outline of the talk

- In the first section I will review essentials of theory of Grothendieck and Street fibrations of categories. They will be the base case of inductive definition of n -fibration in the last section.
- In the second section I will mention how to generalize from Grothendieck fibration to 2-fibration of 2-categories and bicategories.
- The third section will be about internalization of Grothendieck and Street fibrations in 2-categories. Here, I will talk about some very recent work Steve and I did on relating internal fibrations to 2-fibrations.

Outline of the talk

- In the first section I will review essentials of theory of Grothendieck and Street fibrations of categories. They will be the base case of inductive definition of n -fibration in the last section.
- In the second section I will mention how to generalize from Grothendieck fibration to 2-fibration of 2-categories and bicategories.
- The third section will be about internalization of Grothendieck and Street fibrations in 2-categories. Here, I will talk about some very recent work Steve and I did on relating internal fibrations to 2-fibrations.
- In the last section, I will present state of the art regarding theory of n -fibrations of n -categories.

A notation guide

I will stick to these more or less!

- For categories: $\mathcal{A}, \mathcal{B}, \mathcal{C}, \dots, \mathcal{X}$
- Objects of categories, 2-categories, bicategories: A, B, \dots, Z
- For functors: p, q, \dots
- For bicategories: $\mathbb{C}, \mathbb{K}, \mathbb{L}, \dots, \mathbb{X}$
- For 2-functors and bifunctors: $\mathbb{P}, \mathbf{Cod}, \dots$
- Comma categories: $\mathcal{C}^{\rightarrow}, \mathcal{C}^{[1]}, \dots$

Fibrations of groupoids

DEFINITION

A functor $\pi : \mathcal{G} \rightarrow \mathcal{H}$ of groupoids is a **fibration** whenever for every arrow $f : V \rightarrow U$ in \mathcal{H} and every object X in \mathcal{G} sitting above U , there is an arrow $F : Y \rightarrow X$ with $\pi(F) = f$. We call F *lift* of f ending at X .

Fibrations of groupoids

DEFINITION

A functor $\pi : \mathcal{G} \rightarrow \mathcal{H}$ of groupoids is a **fibration** whenever for every arrow $f : V \rightarrow U$ in \mathcal{H} and every object X in \mathcal{G} sitting above U , there is an arrow $F : Y \rightarrow X$ with $\pi(F) = f$. We call F *lift* of f ending at X .

$$\begin{array}{ccc} Y & \xrightarrow{F} & X \\ & \downarrow P & \\ V & \xrightarrow{f} & U \end{array}$$

Fibrations of groupoids

DEFINITION

A functor $\pi : \mathcal{G} \rightarrow \mathcal{H}$ of groupoids is a **fibration** whenever for every arrow $f : V \rightarrow U$ in \mathcal{H} and every object X in \mathcal{G} sitting above U , there is an arrow $F : Y \rightarrow X$ with $\pi(F) = f$. We call F *lift* of f ending at X .

$$\begin{array}{ccc} Y & \xrightarrow{F} & X \\ & \downarrow P & \\ V & \xrightarrow{f} & U \end{array}$$

We call a fibration of groupoids **discrete** whenever all such lifts are unique.

Fibrations of groupoids

DEFINITION

A functor $\pi : \mathcal{G} \rightarrow \mathcal{H}$ of groupoids is a **fibration** whenever for every arrow $f : V \rightarrow U$ in \mathcal{H} and every object X in \mathcal{G} sitting above U , there is an arrow $F : Y \rightarrow X$ with $\pi(F) = f$. We call F *lift* of f ending at X .

$$\begin{array}{ccc} Y & \xrightarrow{F} & X \\ & \downarrow P & \\ V & \xrightarrow{f} & U \end{array}$$

We call a fibration of groupoids **discrete** whenever all such lifts are unique.

EXAMPLE

for a covering map $p : E \rightarrow B$ of topological spaces the fundamental groupoid functor $\Pi(p) : \Pi(E) \rightarrow \Pi(B)$ is a discrete fibration.

Cartesian morphisms

DEFINITION


Suppose \mathcal{X} and \mathcal{C} are categories and $p : \mathcal{X} \rightarrow \mathcal{C}$ is a functor. A morphism $u : Y \rightarrow X$ in \mathcal{X} is called **p-precartesian** whenever for any \mathcal{X} -morphism $v : Z \rightarrow X$ with $p(v) = p(u)$, there exists a unique vertical morphism w such that $u \circ w = v$. Morphism $u : X \rightarrow Y$ is said to be **p-cartesian** whenever for any \mathcal{X} -morphism $v : Z \rightarrow X$ and any $h : p(Z) \rightarrow p(X)$ with $p(u) \circ h = p(v)$, there exists a unique lift w of h such that $u \circ w = v$.

Cartesian morphisms


DEFINITION

Suppose \mathcal{X} and \mathcal{C} are categories and $p : \mathcal{X} \rightarrow \mathcal{C}$ is a functor. A morphism $u : Y \rightarrow X$ in \mathcal{X} is called **p-precartesian** whenever for any \mathcal{X} -morphism $v : Z \rightarrow X$ with $p(v) = p(u)$, there exists a unique vertical morphism w such that $u \circ w = v$. Morphism $u : X \rightarrow Y$ is said to be **p-cartesian** whenever for any \mathcal{X} -morphism $v : Z \rightarrow X$ and any $h : p(Z) \rightarrow p(X)$ with $p(u) \circ h = p(v)$, there exists a unique lift w of h such that $u \circ w = v$.

Precartesian:



Cartesian:



Basic well-known facts about cartesian morphisms

which should not be questioned about in this talk!

- Any cartesian morphism is precartesian.

Basic well-known facts about cartesian morphisms

which should not be questioned about in this talk!

- Any cartesian morphism is precartesian.
- (Pre)cartesian lifts, if they exists, are unique up to unique isomorphism.

Basic well-known facts about cartesian morphisms

which should not be questioned about in this talk!

- Any cartesian morphism is precartesian.
- (Pre)cartesian lifts, if they exist, are unique up to unique isomorphism.
- Any isomorphism is cartesian.

Basic well-known facts about cartesian morphisms

which should not be questioned about in this talk!

- Any cartesian morphism is precartesian.
- (Pre)cartesian lifts, if they exist, are unique up to unique isomorphism.
- Any isomorphism is cartesian.
- Any precartesian vertical arrow in \mathcal{X} is an isomorphism.

Basic well-known facts about cartesian morphisms

which should not be questioned about in this talk!

- Any cartesian morphism is precartesian.
- (Pre)cartesian lifts, if they exist, are unique up to unique isomorphism.
- Any isomorphism is cartesian.
- Any precartesian vertical arrow in \mathcal{X} is an isomorphism.
- Let $v: Y \rightarrow W$ be a \mathfrak{p} -cartesian morphism in \mathcal{X} . Any morphism $u: X \rightarrow Y$ is \mathfrak{p} -cartesian if and only if $v \circ u: X \rightarrow W$ is \mathfrak{p} -cartesian.

Basic well-known facts about cartesian morphisms

which should not be questioned about in this talk!

- Any cartesian morphism is precartesian.
- (Pre)cartesian lifts, if they exist, are unique up to unique isomorphism.
- Any isomorphism is cartesian.
- Any precartesian vertical arrow in \mathcal{X} is an isomorphism.
- Let $v: Y \rightarrow W$ be a \mathfrak{p} -cartesian morphism in \mathcal{X} . Any morphism $u: X \rightarrow Y$ is \mathfrak{p} -cartesian if and only if $v \circ u: X \rightarrow W$ is \mathfrak{p} -cartesian.
- An \mathcal{X} -morphism $u: X \rightarrow Y$ is \mathfrak{p} -cartesian if and only if the following commuting square is a pullback diagram in **Set** for each object Z in \mathcal{X} :

$$\begin{array}{ccc}
 \mathcal{X}(Z, X) & \xrightarrow{u \circ -} & \mathcal{X}(Z, Y) \\
 \mathfrak{p}_{Z, X} \downarrow & \lrcorner & \downarrow \mathfrak{p}_{Z, Y} \\
 \mathcal{C}(\mathfrak{p}Z, \mathfrak{p}X) & \xrightarrow{\mathfrak{p}(u) \circ -} & \mathcal{C}(\mathfrak{p}Z, \mathfrak{p}Y)
 \end{array}$$

Typical example

- For any category \mathcal{C} , there is a codomain functor $\text{cod} : \mathcal{C}^{[1]} \rightarrow \mathcal{C}$ which sends an object $f : J \rightarrow I$ of $\mathcal{C}^{[1]}$ to its codomain C and sends a morphism $\langle v, u \rangle : g \rightarrow f$ of $\mathcal{C}^{[1]}$, i.e. a commuting square, to f .

Typical example

- For any category \mathcal{C} , there is a codomain functor $\text{cod} : \mathcal{C}^{[1]} \rightarrow \mathcal{C}$ which sends an object $f : J \rightarrow I$ of $\mathcal{C}^{[1]}$ to its codomain I and sends a morphism $\langle v, u \rangle : g \rightarrow f$ of $\mathcal{C}^{[1]}$, i.e. a commuting square, to f .
- Cartesian morphisms in $\mathcal{C}^{[1]}$ are precisely pullback squares in \mathcal{C} .

$$\begin{array}{ccc}
 Y & \overset{u^*f}{\dashrightarrow} & X \\
 \lrcorner & & \downarrow u \\
 f^*u & & \\
 \downarrow & & \downarrow f \\
 J & \longrightarrow & I \\
 & & \downarrow \text{cod} \\
 & & J \longrightarrow I \\
 & & f
 \end{array}$$

Fibrations of categories

DEFINITION

A functor $p : \mathcal{X} \rightarrow \mathcal{C}$ is a *Grothendieck pre-fibration* (resp. *Grothendieck fibration*) whenever for each $X \in \mathcal{X}$, every morphism $A \xrightarrow{f} pX$ in \mathcal{C} has a precartesian (resp. cartesian) lift in \mathcal{X} .

Fibrations of categories

DEFINITION

A functor $p : \mathcal{X} \rightarrow \mathcal{C}$ is a *Grothendieck prefibration* (resp. *Grothendieck fibration*) whenever for each $X \in \mathcal{X}$, every morphism $A \xrightarrow{f} pX$ in \mathcal{C} has a precartesian (resp. cartesian) lift in \mathcal{X} .

EXAMPLE

$\text{cod} : \mathcal{C}^{[1]} \rightarrow \mathcal{C}$ is a fibration iff \mathcal{C} has all pullbacks.

Fibrations of categories

DEFINITION

A functor $p : \mathcal{X} \rightarrow \mathcal{C}$ is a *Grothendieck pre-fibration* (resp. *Grothendieck fibration*) whenever for each $X \in \mathcal{X}$, every morphism $A \xrightarrow{f} pX$ in \mathcal{C} has a precartesian (resp. cartesian) lift in \mathcal{X} .

EXAMPLE

$\text{cod} : \mathcal{C}^{[1]} \rightarrow \mathcal{C}$ is a fibration iff \mathcal{C} has all pullbacks.

NON-EXAMPLE

A simple functor which fails to be a fibration: consider category \mathcal{C} consisting of two objects x_0 and x_1 with their identities and an arrow θ_x between them. Let \mathcal{X} be the category extended by a fresh arrow $e : x_0 \rightarrow x_0$ with $e \circ e = e$ and $\theta_x \circ e = \theta_x$. The functor $\mathcal{X} \rightarrow \mathcal{C}$ which sends θ_x to itself and e to id_{x_0} is not a fibration since θ_x in \mathcal{C} does not have a cartesian lift.

More examples of fibrations

- For a category \mathcal{B} , the category of families in \mathcal{B} can be constructed as a comma object in $\mathcal{C}at$. The projection functor $\pi_1: \mathbf{Fam}(\mathcal{B}) \rightarrow \mathbf{Set}$ is a Grothendieck fibration.

$$\begin{array}{ccc} \mathbf{Fam}(\mathcal{B}) & \xrightarrow{!} & \mathbf{1} \\ \pi_1 \downarrow & \nearrow & \downarrow \mathcal{B} \\ \mathbf{Set} & \hookrightarrow & \mathcal{C}at \end{array}$$

More examples of fibrations

- For a category \mathcal{B} , the category of families in \mathcal{B} can be constructed as a comma object in \mathcal{Cat} . The projection functor $\pi_1: \mathbf{Fam}(\mathcal{B}) \rightarrow \mathbf{Set}$ is a Grothendieck fibration.

$$\begin{array}{ccc} \mathbf{Fam}(\mathcal{B}) & \xrightarrow{!} & \mathbf{1} \\ \pi_1 \downarrow & \nearrow & \downarrow \mathcal{B} \\ \mathbf{Set} & \longleftrightarrow & \mathcal{Cat} \end{array}$$

- Wait for a proof of this at the end of this section!

Cloven fibrations

DEFINITION

A **cleavage** for a (pre)fibration $p : \mathcal{X} \rightarrow \mathcal{C}$ is a choice for each $X \in \mathcal{X}$ and $f : B \rightarrow pX$ in \mathcal{C} , a (pre)cartesian lift $\rho(f, X) : \rho_f X \rightarrow X$ of f in \mathcal{X} . More formally, the data of a cleavage is a term ρ of the following dependent type:

$$\rho : \prod_{B, A: \mathbf{Ob}(\mathcal{C})} \prod_{f: \mathcal{C}(B, A)} \prod_{X: \mathcal{X}_A} \sum_{Y: \mathcal{X}_B} \mathit{Cart}_{\mathcal{X}}(Y, X)$$

where the type $\mathit{Cart}_{\mathcal{X}}(Y, X)$ is type of all cartesian morphisms from Y to X . If the fibration p is equipped with a cleavage ρ , then (p, ρ) is called a **cloven fibration**. The cleavage ρ is said to be *splitting* if for any composable pair of morphisms f, g :

$$\rho(g \circ f, X) = \rho(g, X) \circ \rho(f, \rho_g X)$$

And *normalized* whenever for every object X in \mathcal{X} :

$$\rho(id_{pX}, X) = id_X$$

Example of cloven fibration

A cloven fibration $(\text{cod}, \rho) : \mathcal{C}^{[1]} \rightarrow \mathcal{C}$ is precisely a category \mathcal{C} with a choice of pullbacks.

Basic well-known properties of Grothendieck fibrations

which should not be questioned about in this talk!

- A (cloven) prefibration p is a (cloven) fibration iff p -precartesian morphisms are closed under composition.

Basic well-known properties of Grothendieck fibrations

which should not be questioned about in this talk!

- A (cloven) prefibration p is a (cloven) fibration iff p -precartesian morphisms are closed under composition.
- $\langle p, \rho \rangle: \mathcal{X} \rightarrow \mathcal{C}$ is a cloven Grothendieck fibration if and only if for each object $X \in \mathcal{X}$, the induced functor $p_X: \mathcal{X}/X \rightarrow \mathcal{C}/pX$ has a right adjoint right inverse.

Basic well-known properties of Grothendieck fibrations

which should not be questioned about in this talk!

- A (cloven) prefibration \mathfrak{p} is a (cloven) fibration iff \mathfrak{p} -precartesian morphisms are closed under composition.
- $\langle \mathfrak{p}, \rho \rangle: \mathcal{X} \rightarrow \mathcal{C}$ is a cloven Grothendieck fibration if and only if for each object $X \in \mathcal{X}$, the induced functor $\mathfrak{p}_X: \mathcal{X}/X \rightarrow \mathcal{C}/\mathfrak{p}X$ has a right adjoint right inverse.
- $\langle \mathfrak{p}, \rho \rangle: \mathcal{X} \rightarrow \mathcal{C}$ is a cloven Grothendieck fibration if and only if the canonical functor $\mathcal{X}^{[1]} \rightarrow \mathcal{C}/\mathfrak{p}$ has right adjoint right inverse.

Basic well-known properties of Grothendieck fibrations

which should not be questioned about in this talk!

- A (cloven) prefibration \mathfrak{p} is a (cloven) fibration iff \mathfrak{p} -precartesian morphisms are closed under composition.
- $\langle \mathfrak{p}, \rho \rangle: \mathcal{X} \rightarrow \mathcal{C}$ is a cloven Grothendieck fibration if and only if for each object $X \in \mathcal{X}$, the induced functor $\mathfrak{p}_X: \mathcal{X}/X \rightarrow \mathcal{C}/\mathfrak{p}X$ has a right adjoint right inverse.
- $\langle \mathfrak{p}, \rho \rangle: \mathcal{X} \rightarrow \mathcal{C}$ is a cloven Grothendieck fibration if and only if the canonical functor $\mathcal{X}^{[1]} \rightarrow \mathcal{C}/\mathfrak{p}$ has right adjoint right inverse.
- (Cloven) Grothendieck fibrations are closed under composition.

Basic well-known properties of Grothendieck fibrations

which should not be questioned about in this talk!

- A (cloven) prefibration \mathfrak{p} is a (cloven) fibration iff \mathfrak{p} -precartesian morphisms are closed under composition.
- $\langle \mathfrak{p}, \rho \rangle: \mathcal{X} \rightarrow \mathcal{C}$ is a cloven Grothendieck fibration if and only if for each object $X \in \mathcal{X}$, the induced functor $\mathfrak{p}_X: \mathcal{X}/X \rightarrow \mathcal{C}/\mathfrak{p}X$ has a right adjoint right inverse.
- $\langle \mathfrak{p}, \rho \rangle: \mathcal{X} \rightarrow \mathcal{C}$ is a cloven Grothendieck fibration if and only if the canonical functor $\mathcal{X}^{[1]} \rightarrow \mathcal{C}/\mathfrak{p}$ has right adjoint right inverse.
- (Cloven) Grothendieck fibrations are closed under composition.
- (Cloven) Grothendieck fibrations are closed under pullback along other functor.

Basic well-known properties of Grothendieck fibrations

which should not be questioned about in this talk!

- A (cloven) prefibration p is a (cloven) fibration iff p -precartesian morphisms are closed under composition.
- $\langle p, \rho \rangle: \mathcal{X} \rightarrow \mathcal{C}$ is a cloven Grothendieck fibration if and only if for each object $X \in \mathcal{X}$, the induced functor $p_X: \mathcal{X}/X \rightarrow \mathcal{C}/pX$ has a right adjoint right inverse.
- $\langle p, \rho \rangle: \mathcal{X} \rightarrow \mathcal{C}$ is a cloven Grothendieck fibration if and only if the canonical functor $\mathcal{X}^{[1]} \rightarrow \mathcal{C}/p$ has right adjoint right inverse.
- (Cloven) Grothendieck fibrations are closed under composition.
- (Cloven) Grothendieck fibrations are closed under pullback along other functor.
- A Pseudo-pullback of functors is equivalent to their strict pullback if either of the functors is a Grothendieck fibration.

A 2-category of Grothendieck fibrations

DEFINITION

A **(pre)fibration map** between two (pre)fibrations $q : \mathcal{Y} \rightarrow \mathcal{D}$ and $p : \mathcal{X} \rightarrow \mathcal{C}$ consists of two functors $F : \mathcal{D} \rightarrow \mathcal{C}$ and $G : \mathcal{Y} \rightarrow \mathcal{X}$ such that

$$\begin{array}{ccc} \mathcal{Y} & \xrightarrow{G} & \mathcal{X} \\ q \downarrow & & \downarrow p \\ \mathcal{D} & \xrightarrow{F} & \mathcal{C} \end{array}$$

commutes, and moreover, G is cartesian, that is it carries q -cartesian (resp. precartesian) morphisms to p -cartesian (resp. precartesian) morphisms. A **(pre) fibration transformation** is a pair of natural transformations $(\alpha : F_0 \rightarrow F_1, \beta : G_0 \rightarrow G_1)$ such that $p \cdot \beta = \alpha \cdot q$.

A 2-category of Grothendieck fibrations

DEFINITION

A **(pre)fibration map** between two (pre)fibrations $q : \mathcal{Y} \rightarrow \mathcal{D}$ and $p : \mathcal{X} \rightarrow \mathcal{C}$ consists of two functors $F : \mathcal{D} \rightarrow \mathcal{C}$ and $G : \mathcal{Y} \rightarrow \mathcal{X}$ such that

$$\begin{array}{ccc} \mathcal{Y} & \xrightarrow{G} & \mathcal{X} \\ q \downarrow & & \downarrow p \\ \mathcal{D} & \xrightarrow{F} & \mathcal{C} \end{array}$$

commutes, and moreover, G is cartesian, that is it carries q -cartesian (resp. precartesian) morphisms to p -cartesian (resp. precartesian) morphisms. A **(pre) fibration transformation** is a pair of natural transformations $(\alpha : F_0 \rightarrow F_1, \beta : G_0 \rightarrow G_1)$ such that $p \cdot \beta = \alpha \cdot q$.


(Pre)fibrations, (pre)fibration maps, and (pre)fibration transformations form a 2-category.

A 2-category of Grothendieck fibrations


By looking at fibres of a cloven fibration (prefibration) we get a pseudofunctor (resp. lax functor) to 2-category of categories. Grothendieck construction makes this into a biequivalence of 2-categories:

A 2-category of Grothendieck fibrations

By looking at fibres of a cloven fibration (prefibration) we get a pseudofunctor (resp. lax functor) to 2-category of categories. Grothendieck construction makes this into a biequivalence of 2-categories:



A 2-category of Grothendieck fibrations




$$\mathbf{clPreFib}(\mathcal{C}) \simeq \mathbf{LaxFun}(\mathcal{C}^{\text{op}}, \mathcal{Cat})$$

$$\mathbf{clFib}(\mathcal{C}) \simeq \mathbf{PsFun}(\mathcal{C}^{\text{op}}, \mathcal{Cat})$$

where **clFib** is the 2-category of cloven fibrations.

A 2-category of Grothendieck fibrations



REMARK

An interesting feature of the Grothendieck construction is that it reduces category level. That is it turns a 2-functor of 2-categories into a single morphisms in 2-category of categories. Other than a change in viewpoint it makes a world of difference when we work in higher levels; An n -stack in algebraic geometry can be conceived as a category fibred in spaces instead of an ∞ -functor to the ∞ -category \mathcal{S} of spaces.

Fam as example of Grothendieck construction

For a category \mathcal{B} , the Grothendieck fibration $\mathbf{Fam}(\mathcal{B}) \rightarrow \mathbf{Set}$ is the Grothendieck construction of 2-functor

$$\mathbf{Fun}(-, \mathcal{B}): \mathbf{Set}^{\mathrm{op}} \rightarrow \mathcal{C}at$$

where for an (indexing) set I , $\mathbf{Fun}(I, \mathcal{B})$ is the category of functors from discrete category I to category \mathcal{B} .

How forgetful are fibrations?

Homotopy type of homotopy fibre (aka bipullback) over a point tells us how forgetful the forgetful functor/ fibration is. For instance

$$\begin{array}{ccc}
 U^*G & \longrightarrow & \mathbf{Ab} \\
 \downarrow & \cong & \downarrow U \\
 1 & \xrightarrow{G} & \mathbf{Grp}
 \end{array}$$

And one can easily see that the fibre

$$U^*G \cong \left(\sum_{A: \mathbf{Ab}} A \cong G \right)$$

is either empty or a contractible groupoid and in terminology of (Baez and Shulman, 2010), U forgets property of being abelian. However in the case of forgetful fibration $\mathbf{Ab} \rightarrow \mathbf{Set}$ the homotopy fibres are 0-groupoids (aka sets), and the forgetful functor forgets structure of abelian group.

Going one dimension higher

2-cartesian 1-cells

Suppose $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ is a 2-functor. Inspired by the case of 1-functors we define 2-cartesian 1-cells as follows.

Going one dimension higher

2-cartesian 1-cells

Suppose $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ is a 2-functor. Inspired by the case of 1-functors we define 2-cartesian 1-cells as follows.

DEFINITION

A 1-cell $u: X \rightarrow Y$ in \mathbb{X} is **cartesian** with respect to \mathbb{P} whenever for each 0-cell W in \mathbb{X} the following commuting square is a (strict) pullback diagram in 2-category $\mathcal{C}at$.

$$\begin{array}{ccc}
 \mathbb{X}(W, X) & \xrightarrow{u_*} & \mathbb{X}(W, Y) \\
 \mathbb{P}_{W, X} \downarrow & \lrcorner & \downarrow \mathbb{P}_{W, Y} \\
 \mathbb{C}(\mathbb{P}W, \mathbb{P}X) & \xrightarrow{\mathbb{P}(u)_*} & \mathbb{C}(\mathbb{P}W, \mathbb{P}Y)
 \end{array}$$

Going one dimension higher

2-cartesian 1-cells

Suppose $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ is a 2-functor. Inspired by the case of 1-functors we define 2-cartesian 1-cells as follows.

DEFINITION

A 1-cell $u: X \rightarrow Y$ in \mathbb{X} is **cartesian** with respect to \mathbb{P} whenever for each 0-cell W in \mathbb{X} the following commuting square is a (strict) pullback diagram in 2-category $\mathcal{C}at$.


$$\begin{array}{ccc}
 \mathbb{X}(W, X) & \xrightarrow{u_*} & \mathbb{X}(W, Y) \\
 \mathbb{P}_{W,X} \downarrow & \lrcorner & \downarrow \mathbb{P}_{W,Y} \\
 \mathbb{C}(\mathbb{P}W, \mathbb{P}X) & \xrightarrow{\mathbb{P}(u)_*} & \mathbb{C}(\mathbb{P}W, \mathbb{P}Y)
 \end{array}$$

REMARK

By considering object component of pullback diagram above we observe that every 2-cartesian 1-cell is automatically 1-cartesian in the usual sense.

2-cartesian 1-cells in elementary terms

This definition gives us two layers of cartesian properties of 1-cells w.r.t. \mathbb{P} in \mathbb{X} . First of all, u is 1-cartesian as usual. Second, every 2-cell $\alpha: v \Rightarrow w: W \rightarrow Y$ and every 2-cell $\sigma: g \Rightarrow h: \mathbb{P}W \rightarrow \mathbb{P}X$ with $\mathbb{P}(\alpha) = \mathbb{P}(u) \cdot \sigma$ there is a unique lift ϕ of σ such that $u \cdot \phi = \alpha$.



Strict 2-fibrations

DEFINITION

A 2-functor $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ is a **2-fibration** if

- 1 any 1-cell in \mathbb{C} of the form $f: A \rightarrow \mathbb{P}X$ has a 2-cartesian lift,

Strict 2-fibrations

DEFINITION

A 2-functor $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ is a **2-fibration** if

- 1 any 1-cell in \mathbb{C} of the form $f: A \rightarrow \mathbb{P}X$ has a 2-cartesian lift,
- 2 \mathbb{P} is a local fibration, that is for any pair of objects X, Y in \mathbb{X} , the functor $\mathbb{P}_{X,Y}: \mathbb{X}(X, Y) \rightarrow \mathbb{C}(\mathbb{P}X, \mathbb{P}Y)$ is a Grothendieck fibration,

Strict 2-fibrations

DEFINITION

A 2-functor $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ is a **2-fibration** if

- 1 any 1-cell in \mathbb{C} of the form $f: A \rightarrow \mathbb{P}X$ has a 2-cartesian lift,
- 2 \mathbb{P} is a local fibration, that is for any pair of objects X, Y in \mathbb{X} , the functor $\mathbb{P}_{X,Y}: \mathbb{X}(X, Y) \rightarrow \mathbb{C}(\mathbb{P}X, \mathbb{P}Y)$ is a Grothendieck fibration,
- 3 cartesian 2-cells in \mathbb{X} are closed under pre-composition and post-composition with arbitrary 1-cells.

Strict 2-fibrations

DEFINITION

A 2-functor $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ is a **2-fibration** if

- 1 any 1-cell in \mathbb{C} of the form $f: A \rightarrow \mathbb{P}X$ has a 2-cartesian lift,
- 2 \mathbb{P} is a local fibration, that is for any pair of objects X, Y in \mathbb{X} , the functor $\mathbb{P}_{X,Y}: \mathbb{X}(X, Y) \rightarrow \mathbb{C}(\mathbb{P}X, \mathbb{P}Y)$ is a Grothendieck fibration,
- 3 cartesian 2-cells in \mathbb{X} are closed under pre-composition and post-composition with arbitrary 1-cells.

Strict 2-fibrations

DEFINITION

A 2-functor $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ is a **2-fibration** if

- 1 any 1-cell in \mathbb{C} of the form $f: A \rightarrow \mathbb{P}X$ has a 2-cartesian lift,
- 2 \mathbb{P} is a local fibration, that is for any pair of objects X, Y in \mathbb{X} , the functor $\mathbb{P}_{X,Y}: \mathbb{X}(X, Y) \rightarrow \mathbb{C}(\mathbb{P}X, \mathbb{P}Y)$ is a Grothendieck fibration,
- 3 cartesian 2-cells in \mathbb{X} are closed under pre-composition and post-composition with arbitrary 1-cells.

REMARK

The second condition is equivalent to say that for any morphism g in \mathbb{X} and a 2-cell $\alpha: f \Rightarrow \mathbb{P}g$, there is a cartesian 2-cell $\sigma: f \Rightarrow g$ with $\mathbb{P}\sigma = \alpha$.

An archetypal example of strict 2-fibration

EXAMPLE

The 2-category **Fib** of Grothendieck fibrations is a 2-fibred over 2-category of categories via the codomain functor $\text{cod}: \mathbf{Fib} \rightarrow \mathbf{Cat}$

An archetypal example of strict 2-fibration

EXAMPLE

The 2-category **Fib** of Grothendieck fibrations is a 2-fibred over 2-category of categories via the codomain functor $\text{cod}: \mathbf{Fib} \rightarrow \mathcal{C}at$

To my knowledge this was first proved and written down explicitly in [Claudio Hermida \(1999\)](#). “Some properties of Fib as a fibred 2-category”. In: vol. 134, pp. 83–109.

An archetypal example of strict 2-fibration

EXAMPLE

The 2-category **Fib** of Grothendieck fibrations is a 2-fibred over 2-category of categories via the codomain functor $\text{cod}: \mathbf{Fib} \rightarrow \mathcal{Cat}$

To my knowledge this was first proved and written down explicitly in [Claudio Hermida \(1999\)](#). “Some properties of \mathbf{Fib} as a fibred 2-category”. In: vol. 134, pp. 83–109.

Notice that every fibre $\mathbf{Fib}(\mathcal{C})$ is a 2-category consisting of

- 0-cells: Fibred categories over \mathcal{C}

An archetypal example of strict 2-fibration

EXAMPLE

The 2-category **Fib** of Grothendieck fibrations is a 2-fibred over 2-category of categories via the codomain functor $\text{cod}: \mathbf{Fib} \rightarrow \mathcal{Cat}$

To my knowledge this was first proved and written down explicitly in [Claudio Hermida \(1999\)](#). “Some properties of **Fib** as a fibred 2-category”. In: vol. 134, pp. 83–109.

Notice that every fibre $\mathbf{Fib}(\mathcal{C})$ is a 2-category consisting of

- 0-cells: Fibred categories over \mathcal{C}
- 1-cells: Cartesian functors between fibred categories over \mathcal{C}

An archetypal example of strict 2-fibration

EXAMPLE

The 2-category **Fib** of Grothendieck fibrations is a 2-fibred over 2-category of categories via the codomain functor $\text{cod}: \mathbf{Fib} \rightarrow \mathcal{Cat}$

To my knowledge this was first proved and written down explicitly in [Claudio Hermida \(1999\)](#). “Some properties of **Fib** as a fibred 2-category”. In: vol. 134, pp. 83–109.

Notice that every fibre $\mathbf{Fib}(\mathcal{C})$ is a 2-category consisting of

- 0-cells: Fibred categories over \mathcal{C}
- 1-cells: Cartesian functors between fibred categories over \mathcal{C}
- 2-cells: Vertical natural transformations over \mathcal{C}

An archetypal example of strict 2-fibration

EXAMPLE

The 2-category **Fib** of Grothendieck fibrations is a 2-fibred over 2-category of categories via the codomain functor $\text{cod}: \mathbf{Fib} \rightarrow \mathcal{Cat}$

To my knowledge this was first proved and written down explicitly in [Claudio Hermida \(1999\)](#). “Some properties of \mathbf{Fib} as a fibred 2-category”. In: vol. 134, pp. 83–109.

Notice that every fibre $\mathbf{Fib}(\mathcal{C})$ is a 2-category consisting of

- 0-cells: Fibred categories over \mathcal{C}
- 1-cells: Cartesian functors between fibred categories over \mathcal{C}
- 2-cells: Vertical natural transformations over \mathcal{C}

An archetypal example of strict 2-fibration

EXAMPLE

The 2-category **Fib** of Grothendieck fibrations is a 2-fibred over 2-category of categories via the codomain functor $\text{cod}: \mathbf{Fib} \rightarrow \mathcal{Cat}$

To my knowledge this was first proved and written down explicitly in [Claudio Hermida \(1999\)](#). “Some properties of **Fib** as a fibred 2-category”. In: vol. 134, pp. 83–109.

Notice that every fibre $\mathbf{Fib}(\mathcal{C})$ is a 2-category consisting of

- 0-cells: Fibred categories over \mathcal{C}
- 1-cells: Cartesian functors between fibred categories over \mathcal{C}
- 2-cells: Vertical natural transformations over \mathcal{C}

Notice that every 2-fibration gives rise to a base change 2-functor. In this case, given a functor $H: \mathcal{C} \rightarrow \mathcal{D}$, we get a 2-functor $H^*: \mathbf{Fib}(\mathcal{D}) \rightarrow \mathbf{Fib}(\mathcal{C})$ which takes a fibred category over \mathcal{D} to its strict pullback along H .

Weak 2-cartesian 1-cells

DEFINITION

Suppose $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ is a 2-functor. A 1-cell $f: X \rightarrow Y$ in \mathbb{X} is **weakly cartesian** with respect to \mathbb{P} whenever for each 0-cell W in \mathbb{X} the following commuting square is a bipullback diagram in 2-category $\mathcal{C}at$ of categories.

$$\begin{array}{ccc}
 \mathbb{X}(W, X) & \xrightarrow{f_*} & \mathbb{X}(W, Y) \\
 \mathbb{P}_{W, X} \downarrow & \lrcorner & \downarrow \mathbb{P}_{W, Y} \\
 \mathcal{C}(\mathbb{P}W, \mathbb{P}X) & \xrightarrow{\mathbb{P}(f)_*} & \mathcal{C}(\mathbb{P}W, \mathbb{P}Y)
 \end{array}$$

Weak 2-cartesian 1-cells

DEFINITION

Suppose $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ is a 2-functor. A 1-cell $f: X \rightarrow Y$ in \mathbb{X} is **weakly cartesian** with respect to \mathbb{P} whenever for each 0-cell W in \mathbb{X} the following commuting square is a bipullback diagram in 2-category $\mathcal{C}at$ of categories.


$$\begin{array}{ccc}
 \mathbb{X}(W, X) & \xrightarrow{f_*} & \mathbb{X}(W, Y) \\
 \mathbb{P}_{W, X} \downarrow & \lrcorner & \downarrow \mathbb{P}_{W, Y} \\
 \mathcal{C}(\mathbb{P}W, \mathbb{P}X) & \xrightarrow{\mathbb{P}(f)_*} & \mathcal{C}(\mathbb{P}W, \mathbb{P}Y)
 \end{array}$$

Igor Bakovic (2012). “Fibrations in tricategories”. In: *93rd Peripatetic Seminar on Sheaves and Logic, University of Cambridge* and


Mitchell Buckley (2014). “Fibred 2-categories and bicategories”. In: vol. 218, pp. 1034–1074

Weak 2-cartesian 1-cells in elementary terms

Only a bit more complicated than last one-lifts up to iso



\mathbb{P}



Input data:

- ① $g: W \rightarrow Y$
- ② $h: \mathbb{P}W \rightarrow \mathbb{P}X$
- ③ iso 2-cell $\alpha: \mathbb{P}(f) \circ h \Rightarrow \mathbb{P}(g)$


Output data:

(not necc. unique)

- ① $\hat{h}: W \rightarrow X$
- ② iso 2-cell $\hat{\alpha}: f\hat{h} \Rightarrow g$
- ③ iso 2-cell $\hat{\beta}: \mathbb{P}(\hat{h}) \Rightarrow h$
- ④ an equality of 2-cells
 $\alpha \circ (\mathbb{P}(f) \cdot \hat{\beta}) = \mathbb{P}(\hat{\alpha}) \circ \Phi_{h,f}$

Weak 2-cartesian 1-cells in elementary terms

Continued



Input data:

- ① $\sigma: g \Rightarrow g': W \rightrightarrows Y$
- ② $\delta: \mathbb{P}h \Rightarrow \mathbb{P}h': \mathbb{P}W \rightrightarrows \mathbb{P}X$
- ③ iso 2-cells
 $\alpha: \mathbb{P}(f) \circ h \Rightarrow \mathbb{P}(g)$
 $\alpha': \mathbb{P}(f) \circ h' \Rightarrow \mathbb{P}(g)$
- ④ an equality of 2-cells
 $\alpha' \circ (\mathbb{P}f \cdot \delta) = \mathbb{P}(\sigma) \circ \alpha$

Output data:

- ① unique $\hat{\delta}: \hat{h} \Rightarrow \hat{h}'$
- ② an equality $\hat{\alpha}' \circ (f \cdot \hat{\delta}) = \sigma \circ \hat{\alpha}$
- ③ an equality $\delta \cdot (\hat{\beta}) = \hat{\beta}' \circ \mathbb{P}\hat{\delta}$

Cartesian 2-cells

DEFINITION


A 2-cell $\alpha: f \Rightarrow g: x \rightarrow y$ in \mathbb{X} is **cartesian** if it is cartesian as a 1-cell for the functor $\mathbb{P}_{xy}: \mathbb{X}(x, y) \rightarrow \mathbb{C}(\mathbb{P}_x, \mathbb{P}_y)$.

Cartesian 2-cells

DEFINITION

A 2-cell $\alpha: f \Rightarrow g: x \rightarrow y$ in \mathbb{X} is **cartesian** if it is cartesian as a 1-cell for the functor $\mathbb{P}_{xy}: \mathbb{X}(x, y) \rightarrow \mathbb{C}(\mathbb{P}x, \mathbb{P}y)$.

In elementary terms it means a 2-cell $\alpha: f_0 \Rightarrow f_1: X \rightrightarrows Y$ is cartesian if for any given 1-cell $e: X \rightarrow Y$ and 2-cell $\beta: e \rightarrow f_1$ with $\mathbb{P}\alpha = \mathbb{P}\beta \circ \gamma$ for some 2-cell γ , then there is a unique 2-cell $\tilde{\gamma}$ over γ such that $\alpha = \beta \circ \tilde{\gamma}$.



Weak 2-fibrations

As in the case of strict 2-fibrations, we say that \mathbb{P} is *locally fibred* when $\mathbb{P}_{XY}: \mathbb{X}(x, y) \rightarrow \mathbb{C}(\mathbb{P}X, \mathbb{P}Y)$ is a fibration for all X, Y in \mathbb{X} .

DEFINITION

Let $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ be a 2-functor. We say that \mathbb{P} is a **fibration** whenever

- 1 for any $X \in \mathbb{X}$ and $f: B \rightarrow \mathbb{P}X$ in \mathbb{C} , there is a 2-cartesian 1-cell $\tilde{f}: \tilde{B} \rightarrow X$ with $\mathbb{P}\tilde{f} = f$;

Weak 2-fibrations

As in the case of strict 2-fibrations, we say that \mathbb{P} is *locally fibred* when $\mathbb{P}_{XY}: \mathbb{X}(x, y) \rightarrow \mathbb{C}(\mathbb{P}X, \mathbb{P}Y)$ is a fibration for all X, Y in \mathbb{X} .

DEFINITION

Let $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ be a 2-functor. We say that \mathbb{P} is a **fibration** whenever

- 1 for any $X \in \mathbb{X}$ and $f: B \rightarrow \mathbb{P}X$ in \mathbb{C} , there is a 2-cartesian 1-cell $\tilde{f}: \tilde{B} \rightarrow X$ with $\mathbb{P}\tilde{f} = f$;
- 2 \mathbb{P} is locally fibred

Weak 2-fibrations

As in the case of strict 2-fibrations, we say that \mathbb{P} is *locally fibred* when $\mathbb{P}_{XY}: \mathbb{X}(x, y) \rightarrow \mathbb{C}(\mathbb{P}X, \mathbb{P}Y)$ is a fibration for all X, Y in \mathbb{X} .

DEFINITION

Let $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ be a 2-functor. We say that \mathbb{P} is a **fibration** whenever

- 1 for any $X \in \mathbb{X}$ and $f: B \rightarrow \mathbb{P}X$ in \mathbb{C} , there is a 2-cartesian 1-cell $\tilde{f}: \tilde{B} \rightarrow X$ with $\mathbb{P}\tilde{f} = f$;
- 2 \mathbb{P} is locally fibred
- 3 The horizontal composite of any two cartesian 2-cells is again cartesian.

Weak 2-fibrations

As in the case of strict 2-fibrations, we say that \mathbb{P} is *locally fibred* when $\mathbb{P}_{XY}: \mathbb{X}(x, y) \rightarrow \mathbb{C}(\mathbb{P}X, \mathbb{P}Y)$ is a fibration for all X, Y in \mathbb{X} .

DEFINITION

Let $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ be a 2-functor. We say that \mathbb{P} is a **fibration** whenever

- 1 for any $X \in \mathbb{X}$ and $f: B \rightarrow \mathbb{P}X$ in \mathbb{C} , there is a 2-cartesian 1-cell $\tilde{f}: \tilde{B} \rightarrow X$ with $\mathbb{P}\tilde{f} = f$;
- 2 \mathbb{P} is locally fibred
- 3 The horizontal composite of any two cartesian 2-cells is again cartesian.

Weak 2-fibrations

As in the case of strict 2-fibrations, we say that \mathbb{P} is *locally fibred* when $\mathbb{P}_{XY}: \mathbb{X}(x, y) \rightarrow \mathbb{C}(\mathbb{P}X, \mathbb{P}Y)$ is a fibration for all X, Y in \mathbb{X} .

DEFINITION

Let $\mathbb{P}: \mathbb{X} \rightarrow \mathbb{C}$ be a 2-functor. We say that \mathbb{P} is a **fibration** whenever

- 1 for any $X \in \mathbb{X}$ and $f: B \rightarrow \mathbb{P}X$ in \mathbb{C} , there is a 2-cartesian 1-cell $\tilde{f}: \tilde{B} \rightarrow X$ with $\mathbb{P}\tilde{f} = f$;
- 2 \mathbb{P} is locally fibred
- 3 The horizontal composite of any two cartesian 2-cells is again cartesian.

Mitchell Buckley (2014). “Fibred 2-categories and bicategories”. In: vol. 218, pp. 1034–1074

Internal fibrations to 2-categories

The first and the most obvious way to internalize definition of Grothendieck fibration in a 2-category is the representable approach. The second approach was developed by Street in (Street, 1974) who introduced two-sided fibrations in 2-categories, followed by two-sided fibrations in bicategories. These fibrations are defined as algebras over certain 2-monads on 2-categories, or hyperdoctrines on bicategories respectively, and Chevalley's internal characterization of fibrations was obtained as a theorem.

Internal fibrations to 2-categories

The third approach was developed by Johnstone in (Johnstone, [1993](#)) which is closer than Streets definition to the spirit of Grothendiecks original definition. For instance, the base change functors is part of data of definition. Johnstone also established the equivalence of his definition with the representable one.

Internal fibrations to 2-categories

Unlike Street's definition, Johnstone's definition does not require strictness of the 2-category nor the existence of the structure of strict pullbacks and comma objects. Indeed, this definition is most suitable for weak 2-categories such as 2-category of toposes where we do not expect diagrams of 1-cells to commute strictly. This definition is also very flexible in terms of existence of bipullbacks: one only needs existence of bipullbacks of the class of 1-cells one would like to define as (op)fibrations. We will call these 1-cells *carrable*.

Internal fibrations to 2-categories

Unlike Street's definition, Johnstone's definition does not require strictness of the 2-category nor the existence of the structure of strict pullbacks and comma objects. Indeed, this definition is most suitable for weak 2-categories such as 2-category of toposes where we do not expect diagrams of 1-cells to commute strictly. This definition is also very flexible in terms of existence of bipullbacks: one only needs existence of bipullbacks of the class of 1-cells one would like to define as (op)fibrations. We will call these 1-cells *carrable*.

Ross Street (1974). "Fibrations and Yoneda's lemma in a 2-category". In: *Lecture Notes in Math., Springer, Berlin* Vol.420, pp. 104–133

Peter Johnstone (1993). "Fibrations and partial products in a 2-category". In: *Applied Categorical Structures* vol.1

Turning iso 2-cells of \mathbb{K} into 1-cells

Suppose \mathbb{K} is a 2-category and \mathbb{I} is the interval category. We can form a new 2-category $\mathbb{K}^{\mathbb{I}} := \mathbf{Fun}_{ps}(\mathbb{I}, \mathbb{K})$ consisting of (strict) 2-functors, pseudo-natural transformations and modifications between them.

Turning iso 2-cells of \mathbb{K} into 1-cells

Suppose \mathbb{K} is a 2-category and \mathbb{I} is the interval category. We can form a new 2-category $\mathbb{K}^{\mathbb{I}} := \mathbf{Fun}_{ps}(\mathbb{I}, \mathbb{K})$ consisting of (strict) 2-functors, pseudo-natural transformations and modifications between them. More explicitly,

- 0-cells are of the form

$$\begin{array}{c} E \\ p \downarrow \\ S \end{array}$$

where $p \in \mathbb{K}_1$.

Turning iso 2-cells of \mathbb{K} into 1-cells

Suppose \mathbb{K} is a 2-category and \mathbb{I} is the interval category. We can form a new 2-category $\mathbb{K}^{\mathbb{I}} := \mathbf{Fun}_{ps}(\mathbb{I}, \mathbb{K})$ consisting of (strict) 2-functors, pseudo-natural transformations and modifications between them. More explicitly,

- 0-cells are of the form

$$\begin{array}{c} E \\ p \downarrow \\ S \end{array}$$


where $p \in \mathbb{K}_1$.

- 1-cells from q to p are of the form $f = \langle \bar{f}, \overset{\blacktriangledown}{f}, \underline{f} \rangle$

$$\begin{array}{ccc} F & \xrightarrow{\bar{f}} & E \\ q \downarrow & \overset{\blacktriangledown}{f} \Downarrow & \downarrow p \\ A & \xrightarrow{\underline{f}} & S \end{array}$$

where $\overset{\blacktriangledown}{f} : p\bar{f} \Rightarrow \underline{f}q$ is an iso 2-cell in \mathbb{K} .

- 2-cells between 1-cells f and g are of the form $\alpha = \langle \bar{\alpha}, \underline{\alpha} \rangle$ where $\bar{\alpha} : \bar{f} \Rightarrow \bar{g}$ and $\underline{\alpha} : \underline{f} \Rightarrow \underline{g}$ are 2-cells in \mathbb{K}



in such a way that the obvious diagram of 2-cells commutes.

REMARK

\mathbb{K}^I is a globular triple category.

REMARK

\mathbb{K}^I is a globular triple category.

Recall that a triple category is an internal category in the category \mathcal{Dbl} of strict double categories and strict double functors. Also recall that a double category is an internal category in the category \mathcal{Cat} of small categories and functors.

Fibration 0-cells in 2-category $\mathbb{K}^{\mathbb{I}}$

Let $p: E \rightarrow S$ be a carrable 1-cell in \mathbb{K} . We call p a **fibration** 0-cell in 2-category $\mathbb{K}^{\mathbb{I}}$ whenever for any 2-cell $\underline{\alpha}: \underline{f} \Rightarrow \underline{g}: A \rightarrow S$ in \mathbb{K} , we have

- a 1-cell $\langle \overline{r(\alpha)}, r_{\alpha}^{\nabla}, 1_A \rangle: \underline{g}^* p \rightarrow \underline{f}^* p$
- and a 2-cell $(\overline{\alpha}, \underline{\alpha}): f \circ r(\alpha) \Rightarrow g$

[Unpack](#)

Fibration 0-cells in 2-category $\mathbb{K}^{\mathbb{I}}$

Let $p: E \rightarrow S$ be a carrable 1-cell in \mathbb{K} . We call p a **fibration** 0-cell in 2-category $\mathbb{K}^{\mathbb{I}}$ whenever for any 2-cell $\underline{\alpha}: \underline{f} \Rightarrow \underline{g}: A \rightarrow S$ in \mathbb{K} , we have

- a 1-cell $\langle \overline{r(\alpha)}, r_{\alpha}^{\nabla}, 1_A \rangle: \underline{g}^* p \rightarrow \underline{f}^* p$
- and a 2-cell $(\overline{\alpha}, \underline{\alpha}): \underline{f} \circ r(\alpha) \Rightarrow \underline{g}$

in $\mathbb{K}^{\mathbb{I}}$, and moreover the following axioms are satisfied:

- 1 There is an isomorphism $(\overline{\tau_f}, id_{1_A}): id_{\underline{f}^* p} \Rightarrow r_{\underline{f}}$ such that $(\overline{id_f}, id_{\underline{f}}) \circ (\overline{f\tau_0}, id_{\underline{f}}) = (id_{\overline{f}}, id_{\underline{f}})$.

Unpack

Unpack

Fibration 0-cells in 2-category $\mathbb{K}^{\mathbb{I}}$

Let $p: E \rightarrow S$ be a carrable 1-cell in \mathbb{K} . We call p a **fibration** 0-cell in 2-category $\mathbb{K}^{\mathbb{I}}$ whenever for any 2-cell $\underline{\alpha}: \underline{f} \Rightarrow \underline{g}: A \rightarrow S$ in \mathbb{K} , we have

- a 1-cell $\langle \overline{r(\alpha)}, r_{\alpha}^{\nabla}, 1_A \rangle: \underline{g}^* p \rightarrow \underline{f}^* p$
- and a 2-cell $(\overline{\alpha}, \underline{\alpha}): f \circ r(\alpha) \Rightarrow g$

Unpack

in $\mathbb{K}^{\mathbb{I}}$, and moreover the following axioms are satisfied:

- 1 There is an isomorphism $(\overline{\tau_f}, id_{1_A}): id_{\underline{f}^* p} \Rightarrow r_{\underline{f}}$ such that $(\overline{id_f}, id_{\underline{f}}) \circ (\overline{f\tau_0}, id_{\underline{f}}) = (id_{\overline{f}}, id_{\underline{f}})$.

Unpack

- 2 If $\underline{\beta}: \underline{g} \Rightarrow \underline{h}$ is another 2-cell in \mathbb{K} , then there exists an iso 2-cell $\tau_{\alpha, \beta}: r(\alpha) \circ r(\beta) \Rightarrow r(\beta\alpha)$ such that the following diagram of 2-cells in $\mathbb{K}^{\mathbb{I}}$ commutes:

$$\begin{array}{ccc}
 f \circ r(\alpha) \circ r(\beta) & \xrightarrow{\alpha \cdot r(\beta)} & g \circ r(\beta) \\
 f \cdot \tau_{\alpha, \beta} \Downarrow & = & \Downarrow \beta \\
 f \circ r(\beta\alpha) & \xrightarrow{\beta\alpha} & h
 \end{array}$$

- ③ Lifting of α is compatible with left whiskering; That is, given any 1-cell $\underline{k} : B \rightarrow A$ in \mathbb{K} , we require $r(\alpha \cdot k)$ to fit into the following bipullback square in $\mathbb{K}^{\mathbb{I}}$:

$$\begin{array}{ccc}
 (\underline{gk})^* p & \xrightarrow{k_g} & \underline{g}^* p \\
 \downarrow r(\alpha \cdot k) & \cong_{\kappa} & \downarrow r(\alpha) \\
 (\underline{fk})^* p & \xrightarrow{k_f} & \underline{f}^* p
 \end{array}$$

where k_f and k_g are pullback 1-cells over \underline{k} .

We also require pasting of 2-cells α and κ to be equal to 2-cell $\alpha \cdot k$.

Unpack

- ④ For any 1-cells $y = \langle \bar{y}, id, 1_A \rangle$ where $\bar{y}: D \rightarrow \underline{g}^*E$, and $x = \langle \bar{x}, \bar{x}, 1_A \rangle: \underline{g}^*p \circ \bar{y} \rightarrow \underline{f}^*p$ where $\bar{x}: D \rightarrow \underline{f}^*E$, and , any 2-cell $\beta = \langle \bar{\beta}, \underline{\alpha} \rangle: f \circ x \Rightarrow g \circ y$ in $\mathbb{K}^{\mathbb{I}}$ is uniquely factored through α , that is there is a unique 2-cell μ in $\mathbb{K}^{\mathbb{I}}$ with property $(\alpha \cdot y) \circ (f \cdot \mu) = \beta$, that is to say the two pasting diagrams in below are equal:

$$\begin{array}{ccc}
 \underline{g}^*p \circ \bar{y} & \xrightarrow{x} & \underline{f}^*p \\
 y \downarrow & \searrow^{r(\alpha)} & \downarrow f \\
 \underline{g}^*p & \xrightarrow{g} & p
 \end{array}
 \quad \Downarrow \mu
 \quad =
 \quad
 \begin{array}{ccc}
 \underline{g}^*p \circ \bar{y} & \xrightarrow{x} & \underline{f}^*p \\
 y \downarrow & \Downarrow \beta & \downarrow f \\
 \underline{g}^*p & \xrightarrow{g} & p
 \end{array}$$

Unpack

Fibration 0-cells in $\mathbb{K}^{\mathbb{I}}$ are Johnstone fibrations in \mathbb{K}

PROPOSITION

A 0-cell in $\mathbb{K}^{\mathbb{I}}$ is a fibration iff it is a fibration as a 1-cell in \mathbb{K} in the sense of Johnstone.

Fibration 0-cells in $\mathbb{K}^{\mathbb{I}}$ are Johnstone fibrations in \mathbb{K}

PROPOSITION

A 0-cell in $\mathbb{K}^{\mathbb{I}}$ is a fibration iff it is a fibration as a 1-cell in \mathbb{K} in the sense of Johnstone.

EXAMPLE

A 0-cell fibration in 2-category $\mathcal{C}at_{st}^{\mathbb{I}}$ = A Johnstone fibration in 2-category $\mathcal{C}at_{st}$ = A Grothendieck fibration of categories

Fibration 0-cells in $\mathbb{K}^{\mathbb{I}}$ are Johnstone fibrations in \mathbb{K}

PROPOSITION

A 0-cell in $\mathbb{K}^{\mathbb{I}}$ is a fibration iff it is a fibration as a 1-cell in \mathbb{K} in the sense of Johnstone.

EXAMPLE

A 0-cell fibration in 2-category $\mathcal{C}at_{st}^{\mathbb{I}}$ = A Johnstone fibration in 2-category $\mathcal{C}at_{st}$ = A Grothendieck fibration of categories


EXAMPLE

A 0-cell fibration in 2-category $\mathcal{C}at_{ps}^{\mathbb{I}}$ = A Johnstone fibration in 2-category $\mathcal{C}at_{ps}$ = A weak fibration (aka Street fibration) of categories

Fibrations in 2-cats vs. fibrations between 2-cats

PROPOSITION

a 1-cell in $\mathbb{K}^{\mathbb{I}}$ is **Cod**-cartesian iff it is a bipullback square in \mathbb{K} .



Fibrations in 2-cats vs. fibrations between 2-cats

PROPOSITION

A 0-cell p in $\mathbb{K}^{\mathbb{I}}$ is a fibration iff

- Every $\underline{f}: A \rightarrow \mathbf{Cod}(p)$ has a 2-cartesian lift,
- For every 0-cell q , the 2-functor

$$\mathbf{Cod}_{q,p}: \mathbb{K}^{\mathbb{I}}(q, p) \rightarrow \mathbb{K}(\mathbf{Cod}(q), \mathbf{Cod}(p))$$

is a Grothendieck fibration of categories,

- whiskering on the left preserves cartesian 2-cells in $\mathbb{K}^{\mathbb{I}}$.

A few questions to think about

- Weak double categories and more generally n-fold categories are one of the main examples of Segal-type higher categories (Paoli, 2017). Can we generalize the notion of 2-fibration between 2-categories/bicategories to a notion of fibration between weak double categories? For instance, we really like for a closed monoidal category \mathcal{V} , $\mathbf{Dist}(\mathcal{V}) \rightarrow \mathbf{Mon}(\mathcal{V})$ to be a weak double bifibration generalising the Grothendieck bifibration $\mathbf{Mod} \rightarrow \mathbf{Ring}$.

A few questions to think about

- Weak double categories and more generally n-fold categories are one of the main examples of Segal-type higher categories (Paoli, 2017). Can we generalize the notion of 2-fibration between 2-categories/bicategories to a notion of fibration between weak double categories? For instance, we really like for a closed monoidal category \mathcal{V} , $\mathbf{Dist}(\mathcal{V}) \rightarrow \mathbf{Mon}(\mathcal{V})$ to be a weak double bifibration generalising the Grothendieck bifibration $\mathbf{Mod} \rightarrow \mathbf{Ring}$.
- Or further to a notion of fibration between n-fold categories?

A few questions to think about

- Weak double categories and more generally n-fold categories are one of the main examples of Segal-type higher categories (Paoli, 2017). Can we generalize the notion of 2-fibration between 2-categories/bicategories to a notion of fibration between weak double categories? For instance, we really like for a closed monoidal category \mathcal{V} , $\mathbf{Dist}(\mathcal{V}) \rightarrow \mathbf{Mon}(\mathcal{V})$ to be a weak double bifibration generalising the Grothendieck bifibration $\mathbf{Mod} \rightarrow \mathbf{Ring}$.
- Or further to a notion of fibration between n-fold categories?
- Following the first question, since double categories have a well-established notion of weak equivalence, we can ask whether the class of fibrations and weak equivalences form a weak factorization system and whether the fibrant objects are framed bicategories in the sense of (Shulman, 2009)

n -fibrations: a tentative definition

In order to define fibration of (weak) n -categories first we have to say what we mean by a (weak) n -category. There has been a two decades of work on this and it is still not a conclusive matter what an n -category should be. A few of them:

- Batanin Leinster: n -globular set with an action of a suitable globular operad.
- Street: simplicial set satisfying certain horn-filling conditions \simeq truncation of a definition of ω -category
- Tamsamani Simpson: a simplicial object in $(n - 1)$ -categories satisfying object-discreteness and the Segal condition aka a Segal n -category
- Baez- Dolan: an opetopic set having enough n -universal fillers, ...

n -fibrations: a tentative definition

The idea of following tentative definition was put forward initially by Baez and Shulman in [John C. Baez and Michael Shulman \(2010\)](#). “Lectures on n -Categories and Cohomology”. In: *Baez J., May J. (eds) Towards Higher Categories, Springer, New York, NY*. URL: [arXiv:math/0608420](https://arxiv.org/abs/math/0608420) and later more explicitly on nLab by Shulman:

n -fibrations: a tentative definition

The idea of following tentative definition was put forward initially by Baez and Shulman in [John C. Baez and Michael Shulman \(2010\)](#). “Lectures on n -Categories and Cohomology”. In: *Baez J., May J. (eds) Towards Higher Categories, Springer, New York, NY*. URL: [arXiv:math/0608420](#) and later more explicitly on nLab by Shulman:

DEFINITION

Let $P: \mathbb{X} \rightarrow \mathbb{C}$ be a functor between (weak) n -categories. A morphism $f: X \rightarrow Y$ in \mathbb{X} is **cartesian** w.r.t P if for any $W \in \mathbb{X}_0$, the following square:

$$\begin{array}{ccc}
 \mathbb{X}(W, X) & \xrightarrow{f_*} & \mathbb{X}(W, Y) \\
 P_{W,X} \downarrow & \lrcorner & \downarrow P_{W,Y} \\
 \mathbb{C}(PW, PX) & \xrightarrow{P(f)_*} & \mathbb{C}(PW, PY)
 \end{array}$$

is a (weak) pullback of $(n - 1)$ -categories.

n -fibrations: a tentative definition

DEFINITION

We say that $P: \mathbb{X} \rightarrow \mathbb{C}$ is a **weak n -fibration** if

- ① For any object $X \in \mathbb{X}_0$ and morphism $f: X \rightarrow PA$ in \mathbb{C} , there exists a cartesian morphism $\tilde{f}: \tilde{X} \rightarrow A$ and an equivalence $P(\tilde{f}) \simeq f$ in the slice n -category \mathbb{C}/PA ,
- ② For any objects $X, Y \in \mathbb{X}_0$, the functor $P_{X,Y}: \mathbb{X}(X, Y) \rightarrow \mathbb{C}(PX, PY)$ is an $(n-1)$ -fibration,
- ③ For any $W, X, Y \in \mathbb{X}_0$, the square

$$\begin{array}{ccc}
 \mathbb{X}(X, Y) \times \mathbb{X}(W, X) & \xrightarrow{\circ_{\mathbb{X}}} & \mathbb{X}(W, Y) \\
 P_{X,Y} \times P_{W,X} \downarrow & & \downarrow P_{W,Y} \\
 \mathbb{C}(PX, PY) \times \mathbb{C}(PW, PX) & \xrightarrow{\circ_{\mathbb{C}}} & \mathbb{C}(PW, PY)
 \end{array}$$

is a morphism of $(n-1)$ -fibrations.

n -fibrations: a tentative definition

REMARK

An n -fibration is **strict** if in the first condition of definition above, equivalence is replaced with equality. The idea is that a strict 1-fibration should correspond to a Grothendieck fibration of categories while a weak 1-fibration corresponds to a Street fibration of categories.

References

-  Baez, John C. and Michael Shulman (2010). “Lectures on n-Categories and Cohomology”. In: *Baez J., May J. (eds) Towards Higher Categories, Springer, New York, NY*. URL: [arXiv:math/0608420](https://arxiv.org/abs/math/0608420).
-  Bakovic, Igor (2012). “Fibrations in tricategories”. In: *93rd Peripatetic Seminar on Sheaves and Logic, University of Cambridge*.
-  Buckley, Mitchell (2014). “Fibred 2-categories and bicategories”. In: vol. 218, pp. 1034–1074.
-  Hermida, Claudio (1999). “Some properties of Fib as a fibred 2-category”. In: vol. 134, pp. 83–109.
-  Johnstone, Peter (1993). “Fibrations and partial products in a 2-category”. In: *Applied Categorical Structures* vol.1.
-  Paoli, Simona (2017). “Segal-type models of higher categories”. In: 2, p. 298. URL: [arXiv:1707.01868](https://arxiv.org/abs/1707.01868).
-  Shulman, Mike (2009). “Framed bicategories and monoidal fibrations”. In: URL: [arXiv:0706.1286](https://arxiv.org/abs/0706.1286).
-  Street, Ross (1974). “Fibrations and Yoneda’s lemma in a 2-category”. In: *Lecture Notes in Math., Springer, Berlin* Vol.420. pp. 104–133.


End!

THANK YOU FOR YOUR ATTENTION!

Bi-pullback: review

A bi-pullback of an opspan $A \xrightarrow{f} C \xleftarrow{g} B$ in a 2-category \mathbb{K} is given by a 0-cell P together with 1-cells d_0, d_1 and an iso 2-cell $\delta: fd_0 \Rightarrow gd_1$ satisfying a universal property which states that given another iso cone $(l_0, l_1, \lambda: fl_0 \cong gl_1)$ over f, g (with vertex X) there exists a 1-cell u with two iso 2-cells γ_0 and γ_1 such that the pasting diagrams below are equal

Bi-pullback: review



Bi-pullback: review

and furthermore, given 1-cells $u, v: X \rightrightarrows P$ and 2-cells $\alpha: d_0 u \rightrightarrows d_0 v$ and $\beta: d_1 u \rightrightarrows d_1 v$ in such a way that


$$\begin{array}{ccc} fd_0 u & \xrightarrow{f.\alpha} & fd_0 v \\ \delta.u \downarrow & & \downarrow \delta.v \\ gd_1 u & \xrightarrow{g.\beta} & gd_1 v \end{array}$$

commutes, there exists a unique 2-cell $\sigma: u \rightrightarrows v$ such that $d_0 \cdot \sigma = \alpha$ and $d_1 \cdot \sigma = \beta$.

[back to presentation](#)

Supplemental diagrams


Unpacking them yields the following diagram in \mathbb{K} :



where obvious diagram of 2-cells commutes.

Supplemental diagrams

Unpacking τ_f yields the following diagram in \mathbb{K} :



We also get

$$\begin{aligned} \nabla_{\underline{f}} \circ (\underline{f}^* p \cdot \overline{\tau_f}) &= id_{\underline{f}^* p} \\ \overline{id_{\underline{f}}} \circ (\overline{f} \overline{\tau_f}) &= id_{\overline{f}} \end{aligned}$$

Supplemental diagrams


Unpacking $\tau_{\alpha,\beta}$ yields the following diagram in \mathbb{K} :

$$\begin{array}{ccccc}
 & & \overline{r(\beta \circ \alpha)} & & \\
 & & \curvearrowright & & \\
 & \underline{h}^* E & \xrightarrow{\overline{r(\beta)}} & \underline{g}^* E & \xrightarrow{\overline{r(\alpha)}} & \underline{f}^* E \\
 & \downarrow \underline{h}^* p & \cong & \downarrow \underline{g}^* p & \cong & \downarrow \underline{f}^* p \\
 & A & \xrightarrow{1} & A & \xrightarrow{1} & A
 \end{array}$$

Furthermore, we get

$$\begin{aligned}
 \overline{r_{\beta\alpha}} \circ (\overline{f} \cdot \overline{\tau}_{\alpha,\beta}) &= \overline{\beta} \circ (\overline{\alpha} \cdot \overline{r(\beta)}) \\
 r_{\beta\alpha}^\nabla \circ (\underline{f}^* p \cdot \overline{\tau}_{\alpha,\beta}) &= r_\beta^\nabla \circ (r_\alpha^\nabla \cdot r_\beta^\nabla)
 \end{aligned}$$

$\overline{r(\alpha \cdot k)}$ is isomorphic to the bi-pullback of $\overline{r(\alpha)}$ along $\overline{k_f}$, which is to say the top left vertical square of the diagram commutes up to an isomorphism.



Supplemental diagrams

